arXiv:1703.05594v1 [hep-ph] 15 Mar 2017

Black Holes-Hedgehogs and Strings as Defects of the Universal Vacua

B.G. Sidharth [1]*; **C.R. Das** [2]†; **L.V. Laperashvili** [3]‡; **and H.B. Nielsen**[4]§

[1] *International Institute of Applicable Mathematics and Information Sciences, B.M. Birla Science Centre Adarsh Nagar, 500063 Hyderabad, India*

[2] *Bogoliubov Laboratory of Theoretical Physics Joint Institute for Nuclear Research International Intergovernmental Organization, Joliot-Curie 6, 141980 Dubna, Moscow region, Russia*

[3] *The Institute of Theoretical and Experimental Physics, National Research Center "Kurchatov Institute", Bolshaya Cheremushkinskaya, 25, 117218 Moscow, Russia*

[4] *Niels Bohr Institute, Blegdamsvej, 17-21, DK 2100 Copenhagen, Denmark*

Keywords: black holes, hedgehogs, topological defects, multiple point principle, effective potential, cosmological constant, degenerate vacua

PACS: 04.50.Kd, 98.80.Cq, 12.10.-g, 95.35.+d, 95.36.+x

*iiamisbgs@yahoo.co.in, birlasc@gmail.com
†das@theor.jinr.ru
‡laper@itep.ru
§hbech@nbi.dk

Abstract

In the present paper, assuming the Multiple Point Principle (MPP) as a new law of Nature, we considered the existence of the two degenerate vacua in the Standard Model (SM): a) the first Electroweak vacuum at $v = 246$ GeV – "true vacuum", and b) the second Planck scale "false vacuum" at $v_2 \sim 10^{18}$ GeV. In these vacua we investigated topological defects: point defects (monopoles, hedgehogs), line defects (strings, vortices) and sheet defects (bubbles, domain walls). The main aim of this paper is an investigation of the hedgehogs configurations as defects of the false vacuum. In the framework of the $f(R)$ gravity, we considered a black hole solution, which corresponds to a "hedgehog" – global monopole, that has been "swallowed" by a black-hole. Considering the results of the hedgehog lattice theory in the framework of the $SU(2)$ Yang-Mills gauge-invariant theory with hedgehogs in the Wilson loops, we have used the critical value of temperature for the hedgehogs confinement phase. This result gave us the possibility to conclude that hedgehogs can contribute to the SM at the energy scale $\approx 10^4 \sim 10^5$ GeV. Showing a new physics at the scale $\simeq 10 \sim 100$ TeV, hedgehogs can provide the stability of the EW-vacuum and the accuracy of the MPP.

Contents

1 Introduction **4**

2 Degenerate vacua of the Universe **5**

3 Topological defects of the Universal vacua **7**

3.1 Hedgehogs as defects of the false vacuum 8

3.2 Non-differentiable space-time manifold of the vacuum 9

3.3 Hedgehogs in Wilson loops and phase transition in $SU(2)$ Yang-Mills theory 11

4 Higgs mass and vacuum stability/metastability in the Standard Model **13**

5 Hedgehogs as a new physics in the Standard Model **14**

6 Conclusions **15**

1 Introduction

The present paper is devoted to the topological defects of the Universal vacua. During the expansion of the early Universe after the Planck era, different phase transitions resulted in to the formation of the various kind of much discussed topological defects like monopoles or hedgehogs (point defects), strings (line defects) and domain walls (sheet defects). These topological defects appeared due to the breakdown of local or global gauge symmetries.

In this paper we investigated the evolution of bubbles separating two phases: one being the "false vacuum" (Planck scale vacuum) and the other one - the "true vacuum" (EW-scale vacuum). The presence of the hedgehogs, called "defects", is responsible for the destabilization of the false vacuum. Decay of the false vacuum is accompanied by the growth of bubbles of a true vacuum. The defects of the true vacuum are Abrikosov-Nielsen-Olesen vortices (strings) and Sidharth's Compton phase objects. We considered a possibility to have an arbitrary domain wall between these two phases. During the inflation, domain wall annihilates, producing gravitational waves and a lot of SM particles, having masses.

The main idea of this paper is the investigation of the hedgehogs configurations as defects of the false vacuum. We obtained a solution for a black-hole in the region which contains a global monopole in the framework of the $f(R)$ gravity, where $f(R)$ is a function of the Ricci scalar R. We considered the gravitational field, which is an isovector scalar Φ^a with $a = 1, 2, 3$, produced by a spherically symmetric configuration in the scalar field theory with a global $SO(3)$ symmetry. This solution is pointing radially: Φ^a is parallel to $\hat{r}$ – the unit vector in the radial direction, what means that we have a "hedgehog" solution by Alexander Polyakov's terminology. This black-hole solution corresponds to a global monopole that has been "swallowed" by a black-hole.

We have used the results obtained in the framework of $SU(2)$ Yang-Mills theory of the gauge-invariant hedgehog-like structures in the Wilson loops. In particular, we have considered the lattice theory giving the critical value of temperature for the hedgehogs confinement phase. Using the results of the hedgehog lattice theory, we conclude that hedgehogs can contribute to the SM at the energy scale $\approx 10^4 \sim 10^5$ GeV, showing a new physics at the scale $\simeq 10 \sim 100$ TeV. In Section 2 we presented the Multiple Point Principle (MPP) suggested by D.L. Bennett and H.B. Nielsen [4], who postulated the existence in Nature of several degenerate vacua with very small energy density (cosmological constants). It was shown that in the assumption of the existence of the two degenerate vacua in the SM (Electroweak vacuum at $v_1 = v = 246$ GeV, and Planck scale one at $v_2 \sim 10^{18}$ GeV), Froggatt and Nielsen [32] obtained the first prediction of the top-quark and Higgs boson masses, which was further improved by several authors in the next approximations. Section 3 is devoted to the general properties of topological defects of the Universal vacua. We considered topological defects of the "false vacuum" as spherical domain walls (bubbles) or circular loops of cosmic strings, which can be spontaneously produced in a de Sitter-like Universe. The space-time inside the bubble, which we shall refer to as the true vacuum, has the geometry of an open FLRW Universe. In Subsection 3.1 we investigated the hedgehogs configurations as defects of the false vacuum. We obtained a solution for a black-hole in the framework of the $f(R)$ gravity, which corresponds to a global monopole "swallowed" by a black-hole. In Subsection 3.2 we considered a noncommutative description of the space-time manifold of the Universal vacuum, leading to

the almost zero cosmological constant. In Subsection 3.3 hedgehogs in Wilson loops of the $SU(2)$ Yang-Mills theory, and phase transitions in this theory, were investigated using the results of Ref. [57] by V.A. Belavin, M.N. Chernodub and I.E. Kozlov. Their lattice results gave the critical value of the temperature for the hedgehogs confinement phase: $\beta_{crit} \approx 2.5$. In Section 4 we considered the hedgehogs contribution correcting the effective Higgs mass and the problem of the EW-vacuum stability/metastability in the SM. The results of Ref. [57] show that hedgehogs can contribute to the SM at the energy scale $\approx 10^4 \sim 10^5$ GeV, providing the stability of the EW-vacuum. In Section 5 we assume that hedgehogs create a new physics at the scale $\approx 10 \sim 100$ TeV, and this new physics can provide the stability of the EW-vacuum and the validity of the MPP concept.

2 Degenerate vacua of the Universe

This paper is based on the discovery that cosmological constant of our Universe is extremely small, almost zero [1–3]. As a result, we have a new law of Nature which was named as a Multiple Point Principle (MPP) by D.L. Bennett and H.B. Nielsen [4] who first suggested this principle.

MPP means: *There exists in Nature several degenerate vacua with very small energy density, or cosmological constants (CC).* The MPP theory was developed in a lot of papers by H.B. Nielsen, D.L. Bennett, C.D. Froggatt, R.B. Nevzorov, L.V. Laperashvili, C.R. Das (see for example, references [5–22] and recently by other authors [23–26]).

Vacuum energy density of our Universe is the Dark Energy (DE). It is related with cosmological constant (CC) by the following way:

$$\rho_{DE} = \rho_{vac} = (M_{Pl}^{red})^2 \Lambda, \tag{1}$$

where M_{Pl}^{red} is the reduced Planck mass: $M_{Pl}^{red} \simeq 2.43 \times 10^{18}$ GeV. At present, cosmological measurements give:

$$\rho_{DE} \simeq (2 \times 10^{-3} \text{ eV})^4, \tag{2}$$

what means a tiny value of CC:

$$\Lambda \simeq 10^{-84} \text{ GeV}^2. \tag{3}$$

This tiny value of ρ_{DE} was first predicted by B.G. Sidharth in 1997 [1,2]. In 1998 year S. Perlmutter, B. Schmidt and A. Riess [3] were awarded by the Nobel Prize for discovery of the Universe's accelerating expansion.

We see, that we have an extremely small cosmological constant of our Universe. By this reason, Bennett, Froggatt and Nielsen assumed to consider only zero, or almost zero, cosmological constants for all vacua existing in Nature.

Restricted to pure Standard Model (SM) we assumed the existence of only three vacua:

1. **Present Electroweak vacuum**, in which we live.
   It has vacuum expectation value (VEV) of the Higgs field equal to:

$$v_1 = v = \langle \phi_H \rangle \approx 246 \text{ GeV}. \tag{4}$$

2. **High Higgs field vacuum** – Planck scale vacuum, which has the following VEV:

$$v_2 = v = \langle \phi_H \rangle \sim 10^{18} \text{ GeV}. \tag{5}$$

3. **Condensate vacuum.** This third vacuum is a very speculative possible state inside the pure SM, which contains a lot of strongly bound states, each bound from 6 top + 6 anti-top quarks (see Refs. [27–31]).

From experimental results for these three vacua, cosmological constants – minimum of the Higgs effective potentials $V_{eff}(\phi_H)$ – are not exactly equal to zero. Nevertheless, they are extremely small. By this reason, Bennett, Froggatt and Nielsen assumed to consider zero cosmological constants as a good approximation. Then according to the MPP, we have a model of pure SM being finetuned in such a way that these three vacua proposed have just zero energy density.

If the effective potential has three degenerate minima, then the following requirements are satisfied:

$$V_{eff}(\phi^2_{min1}) = V_{eff}(\phi^2_{min2}) = V_{eff}(\phi^2_{min3}) = 0, \tag{6}$$

and

$$V'_{eff}(\phi^2_{min1}) = V'_{eff}(\phi^2_{min2}) = V'_{eff}(\phi^2_{min3}) = 0, \tag{7}$$

where

$$V'(\phi^2) = \frac{\partial V}{\partial \phi^2}. \tag{8}$$

Here we assume that:

$$V_{eff}(\phi^2_{min1}) = V_{present}, \quad V_{eff}(\phi^2_{min2}) = V_{high\ field}, \quad \text{and} \quad V_{eff}(\phi^2_{min3}) = V_{condensate}. \tag{9}$$

As a result, Multiple Point Principle postulates: **There are three vacua in the SM with the same energy density, or cosmological constant, and all cosmological constants are zero, or approximately zero.**

Our MPP is really just an assumption about the coupling constants, i.e. MPP provides three restrictions between the parameters of the SM.

Here we present explicitly the following parameters of the SM:

Λ_{CC} : The cosmological constant; g_t : The top Yukawa coupling;
m_H^2 : Higgs mass squared; Λ_{QCD} : The scale parameter of QCD.

$V_{present}$, $V_{high\ field}$ and $V_{condensate}$ are the vacuum energy densities for the three speculated vacua. Here $V_{present} = 0$ fixes the cosmological constant Λ_{CC} to be essentially zero (it indeed is very small). Then $V_{highfield}$ (meaning the energy density of the vacuum having the very high Higgs field $\phi_H \sim 10^{18}$ GeV) could be taken to predict the Higgs mass. And the $V_{condensate} = 0$ helps to predict the g_t Yukawa coupling. Assuming the existence of the two degenerate vacua in the SM: a) the first Electroweak vacuum at $v_1 = v = 246$ GeV, and b) the second Planck scale vacuum at $v_2 \sim 10^{18}$ GeV, Froggatt and Nielsen predicted in Ref. [32] the top-quark and Higgs boson masses:

$$M_t = 173 \pm 5 \text{ GeV}; \quad M_H = 135 \pm 10 \text{ GeV}. \tag{10}$$

In Fig. 1 it is shown the existence of the second (non-standard) minimum of the effective potential in the pure SM at the Planck scale.

3 Topological defects of the Universal vacua

Topological structures in fields are as important as the fields themselves. The presence of defects determine special features of the vacuum structure.

It is well known that in the early Universe topological defects may be created in the vacuum during the vacuum phase transitions. It is thought that the early Universe underwent a series of phase transitions, each one spontaneously breaking some symmetry in particle physics and giving rise to topological defects of some kind, which in many cases can play an essential role throughout the subsequent evolution of the Universe.

In the context of the General Relativity, Barriola and Vilenkin (see Ref. [33]) studied the gravitational effects of a global monopole as a spherically symmetric topological defect. The authors found, that the gravitational effect of global monopole is repulsive in nature. Thus, one may expect that the global monopole and cosmological constants are connected through their common manifestation as the origin of repulsive gravity. Moreover, both cosmological constant and vacuum expectation value (VEV) are connected while the VEV is connected to the topological defects. All these points lead us to a simple conjecture: There must be a common connection among the cosmological constant, topological defects and the vacuum expectation values (VEVs).

Different phase transitions were resulted during the expansion of the early Universe after the Planck era. They produced the formation of the various kind of topological defects: **point defects** (monopoles, hedgehogs, etc.); **line defects** (strings, vortices), and **sheet defects** (for example, domain walls). The topology of the vacuum manifold dictates the nature of these topological defects, appearing due to the breakdown of local or global gauge symmetries.

In the present paper, we shall discuss another potentially observable manifestation of topological defects. It has been shown in Ref. [34] that topological defects, like spherical domain walls and circular loops of cosmic string, can be spontaneously produced in a de Sitter-like Universe. The initial radii of walls and strings are close to the de Sitter horizon, which corresponds to the Universe radius:

$$R_{un} \simeq R_{de\ Sitter\ horizon} \simeq 10^{28}\ \text{cm}. \tag{11}$$

The bubble, which we shall refer to as the **false vacuum**, to be a de Sitter space with a constant expansion rate H_F . It is convenient to use flat de Sitter coordinates to describe the background of the inflating false vacuum:

$$ds^2 = dt^2 - e^{2H_F t}(dr^2 + r^2 d\Omega^2), \tag{12}$$

where

$$d\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2. \tag{13}$$

The space-time inside the bubble, which we shall refer to as a **true vacuum**, has the geometry of an open FLRW Universe:

$$ds^2 = d\tau^2 - a(\tau)^2(d\xi^2 + \sinh^2\xi d\Omega^2). \tag{14}$$

In the true vacuum constant expansion rate is H_T, which has the meaning of the slow-roll inflation rate inside the bubble at the early stages of its evolution. Cosmological theory

of bubbles was developed in a lot of papers by A. Vilenkin and his collaborators (see for example, Refs. [35, 36]). The physical properties of the defects depend on the embedding vacuum.

3.1 Hedgehogs as defects of the false vacuum

The aim of our present investigation is the consideration of the hedgehog configurations (bubbles) as defects in the false vacuum. Developing the results of Refs. [37–40], we obtained a solution for a black-hole in the region, which contains a global monopole in the framework of the $f(R)$ gravity, where $f(R)$ is a function of the Ricci scalar R. Following to the Gravi-weak unification theory, developed by Ref. [41], we considered the following action near the Planck scale:

$$\begin{aligned} S &= \frac{1}{\kappa^2}\int d^4x \left(\frac{1}{2}|\Phi|^2 R - \Lambda + \frac{1}{2}\mathcal{D}^b_\mu(\Phi^b)^\dagger \mathcal{D}^\mu_a \Phi^a \right. \\ &\left. - \frac{1}{4}\lambda(|\Phi^a|^2 - v_2^2)^2 - \frac{1}{4g_2^2}|F^i_{\mu\nu}|^2 + grav.\ terms\right), \end{aligned} \tag{15}$$

where $\kappa^2 = 8\pi G_N$, G_N is the Newton's gravitational constant, Φ^a is the Higgs triplet field ($a = 1, 2, 3$), λ is the Higgs self-interaction coupling, and v_2 is the vacuum expectation value (VEV) of Φ at the Planck scale:

$$v_2 = \langle \Phi_{min2} \rangle \sim 10^{18}\ \text{GeV}. \tag{16}$$

Here $\mathcal{D}_\mu$ is a covariant derivative:

$$\mathcal{D}^a_\mu = \partial_\mu + i\omega^a_\mu + iA^a_\mu, \tag{17}$$

where ω^a_μ is the gravitational spin-connection, and A^a_μ is adjoint representation of the $SU(2)$ gauge field.

The gravitational field, isovector scalar $\Phi^a = (\Phi^1, \Phi^2, \Phi^3)$, produced by a spherically symmetric "hedgehog" configuration in the scalar field theories with a global $SO(3)$ symmetry, is:

$$\Phi^a = v_2 w(r)\frac{x^a}{r}, \tag{18}$$

where $x^a x^a = r^2$ with $(a = 1, 2, 3)$, and $w(r)$ is some structural function. This solution is pointing radially, what means that Φ^a is parallel to $\hat{r}$ – the unit vector in the radial direction. Therefore, we obtained a "hedgehog" solution (see Refs. [42, 43]). The terminology "hedgehog" was first suggested by Alexander Polyakov [43].

In the flat space the hedgehog's core has the size:

$$\delta \sim \frac{1}{\sqrt{\lambda} v_2}, \tag{19}$$

and the mass:

$$M_{core} \sim \frac{v_2}{\sqrt{\lambda}}, \tag{20}$$

which is:

$$M_{BH} \sim M_{Pl} \sim 10^{-5}\ \text{g}, \tag{21}$$

or

$$M_{BH} \sim 10^{18} \text{ GeV}. \tag{22}$$

This is a black-hole solution, which corresponds to a global monopole that has been "swallowed" by a black-hole.

Now we see, that at the Planck scale the Universe vacuum is described by a non-differentiable space-time: by a foam of black-holes, having lattice-like structure, in which sites are black-holes with "hedgehog" monopoles inside them. This manifold is described by a non-commutative geometry (see Refs. [1, 2, 40, 44–49]), what leads to a tiny value of the cosmological constant.

3.2 Non-differentiable space-time manifold of the vacuum

In Refs. [1, 2] B.G. Sidharth predicted:

a. that a cosmological constant is given by a tiny value:

$$\Lambda \sim H_0^2, \tag{23}$$

where H_0 is the Hubble rate in the early Universe:

$$H_0 \simeq 1.5 \times 10^{-42} \text{ GeV}; \tag{24}$$

b. that a Dark Energy density is very small:

$$\rho_{DE} \simeq 10^{-12} \text{ eV}^4 = 10^{-48} \text{ GeV}^4; \tag{25}$$

c. that a very small DE-density provides an accelerating expansion of our Universe after the Big Bang.

Sidharth proceeded from the following points of view [46] : Modern Quantum Gravity (Loop Quantum Gravity, etc.,) deal with a non-differentiable space-time manifold. In such an approach, there exists a minimal space-time cut off, which leads to the non-commutative geometry, a feature shared by the Fuzzy Space-Time also.

If the space-time is fuzzy, non-differentiable, then it has to be described by a non-commutative geometry with the coordinates obeying the following commutation relations:

$$[dx^\mu, dx^\nu] \approx \beta^{\mu\nu} l^2 \neq 0. \tag{26}$$

Eq. (26) is true for any minimal cut off l.

Previously the following commutation relation was considered by H.S. Snyder [50]:

$$[x, p] = \hbar \left(1 + \left(\frac{l}{\hbar} \right)^2 p^2 \right), \ etc., \tag{27}$$

which shows that effectively 4-momentum p is replaced by

$$p \to p \left(1 + \left(\frac{l}{\hbar}\right)^2 p^2\right)^{-1}. \tag{28}$$

Then the energy-momentum formula now becomes as:

$$E^2 = m^2 + p^2 \left(1 + \left(\frac{l}{\hbar}\right)^2 p^2\right)^{-2}, \tag{29}$$

or

$$E^2 \approx m^2 + p^2 - 2\left(\frac{l}{\hbar}\right)^2 p^4. \tag{30}$$

In such a theory the usual energy momentum dispersion relations are modified [47, 48]. In the above equations l stands for a minimal (fundamental) length, which could be the Planck length λ_{Pl}, or for more generally – Compton wavelength λ_c.

Writing Eq. (30) as

$$E = E' + E'', \tag{31}$$

where E' is the usual (old) expression for energy, and E'' is the new additional term in modification. E'' can be easily verified as $E'' = -m_b c^2$ – for boson fields, and $E'' = +m_f c^2$ – for fermion fields with masses m_b, m_f, respectively. These formulas help to identify the DE density, what was first realized by B.G. Sidharth in Ref. [2].

DE density is the density of the quantum vacuum energy of the Universe. Quantum vacuum, described by Zero Point Fields (ZPF) contributions, is the lowest state of any Quantum Field Theory (QFT), and due to the Heisenberg's principle has an infinite value, which is renormalizable.

As it was pointed out in Refs. [44, 51], the quantum vacuum of the Universe can be a source of cosmic repulsion. However, a difficulty in this approach has been that the value of the cosmological constant turns out to be huge [51], far beyond what is observed by astrophysical measurements. This has been called "the cosmological constant problem" [52]. Using the non-commutative theory of the discrete space-time, B.G. Sidharth predicted in Refs. [2, 44] a tiny value of cosmological constant: $\Lambda \simeq 10^{-84}$ GeV^2.

Global monopole is a heavy object formed as a result of gauge-symmetry breaking in the phase transition of the isoscalar triplet Φ^a system. The black-holes-monopoles-hedgehogs are similar to elementary particles, because a major part of their energy is concentrated in a small region near the monopole core.

In the Guendelman-Rabinowitz theory [37] the authors investigated the evolution of bubbles separating two phases:

1. one being the "false vacuum" (Planck scale vacuum), and
2. the other – the "true vacuum" (EW-scale vacuum).

The presence of hedgehogs, called "defects", is responsible for the destabilization of the false vacuum. Decay of the false vacuum is accompanied by the growth of bubbles of a true vacuum. The defects of the true vacuum are Abrikosov-Nielsen-Olesen vortices (strings) [53, 54] and Sidharth's Compton phase objects [55, 56]. Guendelman and Rabinowitz [37] also allowed a possibility to consider an arbitrary domain wall between two phases. During the inflation domain wall annihilates, producing gravitational waves and a lot of the SM particles, having masses.

3.3 Hedgehogs in Wilson loops and phase transition in $SU(2)$ Yang-Mills theory

The authors of Ref. [57] investigated the gauge-invariant hedgehog-like structures in the Wilson loops of the $SU(2)$ Yang-Mills theory. In this model the triplet Higgs field $\hat{\Phi} = \frac{1}{2}\Phi^a\sigma^a$ vanishes at the center of the monopole $x = x_0$: $\Phi(x_0) = 0$, and has a generic hedgehog structure in the spatial vicinity of this monopole.

In the Yang-Mills theory, a hedgehog structure can be entirely defined in terms of Wilson-loop variables [58]. In general, we consider an untraced Wilson loop, beginning and ending at the point x_0 on the closed loop C:

$$W_C(x_0) = P \exp ig \oint_C dx_\mu \hat{A}_\mu. \tag{32}$$

To improve the analogy with the triplet Higgs field $\hat{\Phi}$, we subtract the singlet part from $W_C(x_0)$:

$$\hat{\Gamma}_C(x_0) = W_C(x_0) - \mathbb{1} \cdot \frac{1}{2} Tr W_C(x_0). \tag{33}$$

This is a traceless adjoint operator similar to the field $\hat{\Phi}$. Associating the triplet part $\hat{\Gamma}_C(x_0)$ of Wilson loop $W_C(x_0)$ with the triplet Higgs field $\hat{\Phi}$, we notice the following property: As the Higgs field vanishes in all points x, belonging to the monopole trajectory, similarly Γ_C vanishes on the hedgehog loop C:

$$W_C \in Z_2 \Leftrightarrow \Gamma_C = 0.$$

With aim to demonstrate that the center-valued Wilson loop variables have something to do with the hedgehog structure, we infinitesimally deform the contour $C \to C + \delta C$ in the vicinity of the point x_0. This shift also relocates the reference point, $x_0 \to x_0 + \delta x$ (see Fig. 2). Without loss of generality, we assume that a tangent vector to the loop C at the point x_0 is aimed in the time direction. A physically distinguishable shift δx should be tangential (spacelike) to the contour C because longitudinal (timelike) shifts of the point x_0 leave the functional Γ_C invariant. Then in general case, the deformed loop $\Gamma_C(x)$ should have a spatial hedgehog-like structure in the vicinity of the zero point x_0. Because the hedgehog loops are closed, the hedgehog charges must be conserved.

In conventional superconductivity [53], Abrikosov vortices are singularities in the superconducting condensate (i.e., in the Cooper-pair field). In the core of the Abrikosov vortices, the superconductivity is broken, and the normal metal state is restored. As temperature increases, the condensate weakens, and nucleation of the vortices due to

thermal fluctuations strengthens. Thus, the higher the temperature is, the density of the (thermal) vortices should be larger. It can be expected in the YM theory that the density of hedgehog loops is also sensitive to the phase transition.

The order parameter of the phase transition is the vacuum expectation value (trace) of the Polyakov line:

$$\hat{L}(x) = P \exp ig \int_0^{1/T} dx_4 A_4(\vec{x}, x_4). \tag{34}$$

VEV is $L = \frac{1}{2} Tr \hat{L}$. Functional $\hat{L}(x)$, also called the thermal Wilson line, is a basic variable in an effective theory, which describes the properties of the finite-temperature phase transition of the system. In the confinement phase, the expectation value of the Polyakov line is zero: $\langle L \rangle = e^{-TF_q} = 0$, indicating that the free energy of a single quark becomes infinite when $F_q \to \infty$. In the deconfinement phase, the Polyakov line has a non-zero expectation value: $\langle L \rangle \neq 0$, and the quarks are no longer confined.

Considering lattice model of the $SU(2)$ Yang-Mills theory, Belavin, Chernodub and Kozlov showed numerically that the density of hedgehogs structures in the thermal Wilson-Polyakov lines is very sensitive to the finite-temperature phase transition. The hedgehog line density behaves like an order parameter: the density is almost independent of the temperature in the confinement phase and changes substantially as the system enters the deconfinement phase. These authors obtained a very important result: $\beta_{crit} \approx 2.5$, what means that the critical temperature corresponding to the hedgehogs confinement is smaller than the Planck scale value.

Indeed, $\beta = 1/g^2 = 1/(4\pi\alpha)$, or $\alpha^{-1} = 4\pi\beta$, what means that the critical value α_{crit} for hedgehogs confinement, obtained by the Belavin-Chernodub-Kozlov lattice investigation [57], has the following value:

$$\alpha_{crit}^{-1} \approx 4\pi \times 2.5 \approx 31.4. \tag{35}$$

It is known that the renormalization group equations (RGEs) give $\alpha_{Pl} \approx 50$ for couplings $SU(3)$- and $SU(2)$- gauge theories [59]. In general case, according to RGEs, we have:

$$\alpha^{-1}(\mu) = \alpha_{Pl}^{-1} + bt, \tag{36}$$

where $t = \ln(\mu/M_{Pl})$. Usually RGE is a function of x: $x = \log_{10} \mu$, and $t = x \ln 10 - \ln M_{Pl} \approx (x - 19) \ln 10$. For $SU(2)$-gauge theory $b \approx (19/12\pi)$ and $\alpha_2^{-1}(M_{Pl}) \approx 48$, then we obtain:

$$\alpha_{crit}^{-1} \approx 31.4 = 48 + 0.504(x_{crit} - 19) \ln 10, \tag{37}$$

what gives:

$$x_{crit} \approx 4 \sim 5.$$

This means that hedgehogs can contribute at LHC (maybe only virtually) at energy:

$$E \simeq 10^4 \sim 10^5 \text{ GeV} = 10 \sim 100 \text{ TeV}.$$

4 Higgs mass and vacuum stability/metastability in the Standard Model

As it was mentioned, assuming the existence of two degenerate vacua in the SM (the first Electroweak vacuum and the second Planck scale one), Froggatt and Nielsen predicted the top-quark and Higgs boson masses: $M_t = 173 \pm 5$ GeV and $M_H = 135 \pm 10$ GeV. Their prediction for the mass of the Higgs boson was improved in Ref. [60] by the calculation of the two-loop radiative corrections to the effective Higgs potential. The prediction of Ref. [60]: $M_H = 129 \pm 2$ GeV provided the possibility of the theoretical explanation of the value $M_H \simeq 125.7$ GeV observed at LHC.

The authors of reference [61] extrapolated the SM parameters up to the high (Planck) energies with full 3-loop NNLO RGE precision. From Degrassi et al. calculation [61], the effective Higgs field potential $V_{eff}(\phi_H)$ has a minimum, which goes slightly under zero, so that the present vacuum is unstable for the experimental Higgs mass 125.09 ± 0.24 GeV, while the value that would have made the second minimum just degenerate with the present vacuum energy density would be rather $m_H \simeq 129.4$ GeV.

A theory of a single scalar field is given by the effective potential $V_{eff}(\phi_c)$ which is a function of the classical field ϕ_c. In the loop expansion V_{eff} is given by

$$V_{eff} = V(0) + \Sigma_{n=1} V^{(n)}, \tag{38}$$

where $V(0)$ is the tree-level potential of the SM:

$$V(0) = -\frac{1}{2} m^2 \phi^2 + \frac{1}{4} \lambda \phi^4. \tag{39}$$

The vast majority of the available experimental data is consistent with the SM predictions. No sign of new physics has been detected. Until now there is no evidence for the existence of any particles other than those of the SM, or bound states composed of other particles. All accelerator physics seems to fit well with the SM, except for neutrino oscillations. These results caused a keen interest in possibility of emergence of new physics only at very high (Planck scale) energies, and generated a great attention to the problem of the vacuum stability: whether the electroweak vacuum is stable, unstable, or metastable. A largely explored scenario assumes that new physics interactions only appear at the Planck scale $M_{Pl} = 1.22 \times 10^{19}$ GeV. According to this scenario, we need the knowledge of the Higgs effective potential $V_{eff}(\phi)$ up to very high values of ϕ.

The loop corrections lead the V_{eff} to values of ϕ, which are much larger than $v_1 \approx 246$ GeV – the location of the electroweak (EW) minimum. The effective Higgs potential develops a new minimum at $v_2 \gg v$. The position of the second minimum depends on the SM parameters, especially on the top and Higgs masses, M_t and M_H. This V_{eff} can be higher or lower than the EW one showing a stable EW vacuum (in the first case), or metastable one (in the second case). The red solid line of Fig. 3 by Degrassi et al. shows the running of the $\lambda_{eff}(\phi)$ for $M_H \simeq 125.7$ GeV and $M_t \simeq 171.43$ GeV, which just corresponds to the stability line, that is, to the stable vacuum. In this case the minimum of the $V_{eff}(\phi)$ exists at the $\phi = \phi_0 \simeq 2.22 \times 10^{18}$ GeV, where according to MPP:

$$\lambda_{eff}(\phi_0) = \beta(\lambda_{eff}(\phi_0)) = 0.$$

But as it was shown in Refs. [60, 61], this case does not correspond to the current experimental values.

In Fig. 3 blue lines (thick and dashed) present the RG evolution of $\lambda(\mu)$ for current experimental values $M_H \simeq 125.7$ GeV and $M_t \simeq 173.34$ GeV. The thick blue line corresponds to the central value of $\alpha_s = 0.1184$ and dashed blue lines correspond to its errors equal to ± 0.0007. We see that absolute stability of the Higgs potential is excluded by at 98% C.L. for $M_H < 126$ GeV. Fig. 3 shows that asymptotically $\lambda(\mu)$ does not reach zero, but approaches to the negative value:

$$\lambda \to -0.01 \pm 0.002, \tag{40}$$

indicating the metastability of the EW vacuum. According to the paper of Ref. [60], the stability line shown in Fig. 3 by the red thick line corresponds to $M_H = 129.4 \pm 1.8$ GeV. We see that the current experimental values of M_H and M_t show the metastability of the present Universe vacuum, what means that MPP law is not exact.

5 Hedgehogs as a new physics in the Standard Model

Can the MPP be exact due to corrections from hedgehogs contributions? We think that it is possible. If we assume that in the region of the second minimum of the effective potential there exists an interaction between hedgehog triplet field Φ^a and Higgs doublet H^α (here $a = 1, 2, 3$ and $\alpha = 1, 2$), then we have such additional terms in the action:

$$\begin{aligned} S &= \frac{1}{\kappa^2}\int d^4x \left(f(R) - \Lambda - m_h^2|\Phi|^2 + \lambda_h(\Phi)|\Phi|^4 - m_H^2|H|^2 \right. \\ &\left. + \lambda_H(H)|H|^4 + \lambda_{hH}(h,H)|\Phi|^2|H|^2 + ... \right). \end{aligned} \tag{41}$$

Now we see, that the main Feynman diagram, correcting the effective Higgs self-interaction coupling constant $\lambda_H \equiv \lambda$, is the diagram of Fig. 4, containing the hedgehog Φ (or $h \equiv \Phi$) in the loop. Now we have:

$$\lambda(\mu) = \frac{G_F}{\sqrt{2}} M_H^2 + \Delta\lambda(\mu) + \delta\lambda(\mu), \tag{42}$$

where the term $\delta\lambda(\mu)$ denotes the loop corrections to the Higgs mass arising from hedgehogs, and the main contribution to $\delta\lambda(\mu)$ is the term λ_S:

$$\delta\lambda(\mu) = \lambda_S + ..., \tag{43}$$

which corresponds to the contribution of the diagram of Fig. 4. It is easy to calculate this contribution by methods presented in Ref. [22]:

$$\lambda_S \approx \frac{1}{16\pi^2} \frac{\lambda_{hH}^2}{(r_h m_h)^4}, \tag{44}$$

where λ_{hH} is a coupling constant of the interaction of hedgehogs h with the SM Higgs fields H (see Eq. (41)). In Eq. (47) parameters m_h and r_h are the mass and radius of hedgehog, respectively. According to Eqs. (19) and (20), we have:

$$m_h = M_{core} \quad \text{and} \quad r_h = \delta, \tag{45}$$

and

$$r_h m_h \simeq \frac{1}{\lambda_h}, \tag{46}$$

where λ_h (equal to λ in Eq. (15)) is a self-interaction coupling constant of hedgehog fields Φ. As a result, we obtain:

$$\lambda_S \simeq \frac{\lambda_{hH}^2 \lambda_h^4}{16\pi^2}. \tag{47}$$

If hedgehog parameters λ_h and λ_{hH} give us $\lambda_S \simeq 0.01$, then the hedgehogs contribution transforms the metastable (blue) curve into the stable (red) curve, and we have exact vacuum stability and exact MPP.

This problem needs the further investigation, together with a very important question: can hedgehog's triplet Higgs field be responsible for the Hybrid inflation of the $SU(2)$-doublet Higgs field H?

The main result of this investigation is an assumption that a new physics in the SM comes at the scale $\approx 10 \sim 100$ TeV, and this new physics is presented by the existence of black-holes-hedgehogs in our Universe. This problem is an aim of LHC and new cosmological theory.

6 Conclusions

1. In this investigation we were based on the discovery that cosmological constant of our Universe is extremely small, almost zero, and assumed a new law of Nature which was named as a Multiple Point Principle (MPP). The MPP postulates: *There are two vacua in the SM with the same energy density, or cosmological constant, and both cosmological constants are zero, or approximately zero.* We considered the existence of the following two degenerate vacua in the SM: a) the first Electroweak vacuum at $v_1 = v = 246$ GeV, and b) the second Planck scale vacuum at $v_2 \sim 10^{18}$ GeV.

2. We investigated topological structure of the Universal vacua, assuming that topological defects, like spherical domain walls and circular loops of cosmic strings, can be spontaneously produced in a de Sitter-like Universe. The initial radii of walls and strings are close to the de Sitter horizon, which corresponds to the Universe radius. The bubble, which we shall refer to as the false vacuum, to be a de Sitter space with its constant expansion rate H_F. The space-time inside the bubble, which we shall refer to as the true vacuum, has the geometry of an open FLRW Universe.

3. The aim of this investigation is the consideration of the hedgehogs configurations as defects in the false vacuum. We considered a solution for a black-hole in the region which contains a global monopole in the framework of the $f(R)$ gravity, where $f(R)$ is a function of the Ricci scalar R. The gravitational field, isovector scalar Φ^a with $a = 1, 2, 3$, produced by a spherically symmetric configuration in the scalar field theory with a global $SO(3)$ symmetry, is pointing radially: Φ^a is parallel to $\hat{r}$ – the unit vector in the radial direction, what means that we obtained a "hedgehog" solution by Alexander Polyakov's terminology. This black-hole solution corresponds to a global monopole that has been "swallowed" by a black-hole.

4. We have shown, that the Planck scale Universe vacuum is described by a non-differentiable space-time: by a foam of black-holes, having lattice-like structure, in which sites are black-holes with the "hedgehog" monopoles inside them. This manifold is described by a non-commutative geometry, what leads to a tiny value of cosmological constant $\Lambda \approx 0$.

5. We investigated the evolution of bubbles separating two phases: one being the "false vacuum" (Planck scale vacuum) and the other - the "true vacuum" (EW-scale vacuum). The presence of hedgehogs, called "defects", is responsible for the destabilization of the false vacuum. Decay of the false vacuum is accompanied by the growth of bubbles of a true vacuum. The defects of the true vacuum are Abrikosov-Nielsen-Olesen vortices (strings) and Sidharth's Compton phase objects. We considered a possibility to have an arbitrary domain wall between these two phases. During the inflation domain wall annihilates, producing gravitational waves and a lot of SM particles, having masses.

6. We have used the results by V.A. Belavin, M.N. Chernodub and I.E. Kozlov, who investigated in the framework of $SU(2)$ Yang-Mills theory the gauge-invariant hedgehog-like structures in the Wilson loops. Their lattice theory gave the critical value of the temperature for the hedgehogs confinement phase: $\beta_{crit} \approx 2.5$. This result means that hedgehog field, the Higgs triplet Φ^a, can contribute to the SM at the energy scale $\approx 10^4 \sim 10^5$ GeV.

7. We suggested to develop an assumption that the hedgehog field Φ^a create a new physics at the scale $\approx 10 \sim 100$ TeV, providing the stability of the EW-vacuum and the accuracy of the MPP. These results essentially depend on hedgehog field parameters: mass, radius and coupling constants λ_h (self-interaction of hedgehogs) and λ_{hH} (the interaction of hedgehogs with the SM Higgs fields H).

Acknowledgments

LVL greatly thanks the B.M. Birla Science Centre (Hyderabad, India) and personally Prof. B.G. Sidharth, for hospitality, collaboration and financial support. HBN wishes to thank the Niels Bohr Institute for the status of professor emeritus and corresponding support. CRD is thankful to Prof. D.I. Kazakov for support.

References

[1] B.G. Sidharth, Proc. of the 8th Marcell Grossmann Meeting on General Relativity, Jerusalem, May 1997, ed. T. Piran, (World Scientic, Singapore), pp. 476-479 (1997).

[2] B.G. Sidharth, Int. J. Mod. Phys. A**13**, 2599 (1998), arXiv:quant-ph/9808031.

[3] S. Perlmutter, B. Schmidt and A. Riess, Nature **391**, 51 (1998).

[4] D.L. Bennett and H.B. Nielsen, Int. J. Mod. Phys. A9, 5155 (1994), arXiv:hep-ph/9311321.

[5] D.L. Bennett, C.D. Froggatt and H.B. Nielsen, in Proceedings of the 27th International Conference on High Energy Physics, Glasgow, Scotland, 1994, eds. P. Bussey and I. Knowles (IOP Publishing Ltd., 1995), p. 557.

[6] D.L. Bennett and H.B. Nielsen, Int. J. Mod. Phys. A**14**, 3313 (1999), arXiv:hep-ph/9607278.

[7] D.L. Bennett and H.B.Nielsen, *The multiple point principle: Realized vacuum in nature is maximally degenerate*, Proceedings of Institute of Mathematics of NAS of Ukraine **50**, Part 2, 629-636 (2004).

[8] D.L. Bennett, *The multiple point principle: Characterization of the possible phases for the SMG*, Bled Workshops, Phys. **10**, no.2, p.9 (2009).

[9] D.L. Bennett, L.V. Laperashvili and H.B. Nielsen, *Finestructure constants at the Planck scale from multiple point principle*, in: Proceedings of the 10th Workshop on *What Comes Beyond the Standard Model*, Bled, Slovenia, 17-27 September, 2007, eds. M. Breskvar et al. (Bled Workshops in Physics, Vol. 8 no. 2, DMFA-Zaloznistvo, Ljubljana, 2007), arXiv:0711.4681.

[10] D.L. Bennett, L.V. Laperashvili and H.B. Nielsen, *Relation between ne structure constants at the Planck scale from multiple point principle*, in: Proceedings of the 9th Workshop on *What Comes Beyond the Standard Model*, Bled, Slovenia, 16-26 September, 2006, eds. M. Breskvar et al. (Bled Workshops in Physics, Vol. 7, no. 2, DMFA-Zaloznistvo, Ljubljana, 2006), arXiv:hep-ph/0612250.

[11] C.D. Froggatt, Larisa Laperashvili, R.B. Nevzorov, H.B. Nielsen and Marc Sher, *Implementation of the multiple point principle in the two-Higgs doublet model of type II*, Phys. Rev. D**73**, 095005 (2006), arXiv:hep-ph/0602054.

[12] C.D. Froggatt, R. Nevzorov and H.B. Nielsen, *Smallness of the cosmological constant and the multiple point principle*, J. Phys. Conf. Ser. **110**, 072012 (2008), arXiv:0708.2907.

[13] C.R.Das and L.V. Laperashvili, *Phase transition in gauge theories, monopoles and the multiple point principle*, Int. J. Mod. Phys. A**20**, 5911 (2005), arXiv:hep-ph/0503138.

[14] C.D. Froggatt, L.V. Laperashvili, R.B. Nevzorov, H.B. Nielsen and M. Sher, *The Two Higgs doublet model and the multiple point principle*, Bled Workshops Phys. **5**, no.2, 28-39 (2004), arXiv:hep-ph/0412333.

[15] C. Froggatt, L. Laperashvili, R. Nevzorov and H.B. Nielsen, *No-scale supergravity and the multiple point principle*, Bled Workshops Phys. **5**, no.2, 17-27 (2004), arXiv:hep-ph/0411273.

[16] L.V. Laperashvili, Phys. Atom. Nucl. **57**, 471 (1994) [Yad.Fiz.] **57**, 501 (1994)].

[17] L.V. Laperashvili and H.B.Nielsen, *Multiple point principle and phase transition in gauge theories*, in: Proceedings to the Workshop: *What Comes Beyond the Standard Models?* , Slovenia, Bled, 29 June – 9 July, 1998, DMFA Zaloznistvo, Ljubljana, 1999, arXiv: hep-ph/9905357.

[18] L.V. Laperashvili, *The Multiple point principle and Higgs bosons*, Phys. Part. Nucl. **36**, S38-S40 (2005), arXiv:hep-ph/0411177.

[19] C.D. Froggatt, L.V. Laperashvili, R.B. Nevzorov and H.B. Nielsen, *Cosmological constant in SUGRA models and the multiple point principle*, Phys. Atom. Nucl. **67**, 582 (2004) [Yad. Fiz. **67**, 601 (2004)], arXiv:hep-ph/0310127.

[20] H.B. Nielsen and A. Kleppe, in: Proceedings to the 16th Workshop on *What Comes Beyond the Standard Models?* , Slovenia, Bled, 14-21 July, 2014, Vol.14, No.2, DMFA Zaloznistvo, Ljubljana, arXiv:1403.14101.

[21] C.R. Das, L.V. Laperashvili, H.B. Nielsen, A. Tureanu and C.D. Froggatt, *Gravi-weak unification and multiple-point principle*, Phys. Atom. Nucl. **78** 3, 440 (2015) [Yad. Fiz. **78** 5, 471 (2015)].

[22] L.V. Laperashvili, H.B. Nielsen and C.R. Das, *New results at LHC confirming the vacuum stability and Multiple Point Principle*, Int. J. Mod. Phys. A**31**, 1650029 (2016), arXiv:1601.03231.

[23] G.E. Volovik, *Coexistence of different vacua in the effective quantum field theory and multiple point principle*, JETP Lett. **79**, 101 (2004) [Pisma Zh. Eksp. Teor. Fiz. **79**, 131 (2004)], arXiv:hep-ph/0309144.

[24] Kiyoharu Kawana, *Multiple Point Principle of the Standard Model with Scalar Singlet Dark Matter and Right Handed Neutrinos*, PTEP 2015, 023B04 (2015), arXiv:1411.2097.

[25] Kiyoharu Kawana, *Multiple Point Principle of the Gauged B-L Model*, arXiv:1504.06707.

[26] Naoyuki Haba, Hiroyuki Ishida, Nobuchika Okada, Yuya Yamaguchi, *Multiple-point principle with a scalar singlet extension of the Standard Model*, PTEP 2017 (2017) no.1, 013B03, arXiv:1608.00087.

[27] C.D. Froggatt and H.B. Nielsen, *Trying to understand the Standard Model parameters.* Invited talk by H.B. Nielsen at the "XXXI ITEP Winter School of Physics", Moscow, Russia, 18-26 February, 2003; Surveys High Energy Phys. **18**, 55-75 (2003); hep-ph/0308144.

[28] C.D. Froggatt, H.B. Nielsen and L.V. Laperashvili, *Hierarchy-problem and a bound state of 6 t and 6 anti-t*, in: Proceedings of Coral Gables Conference on Launching of Belle Epoque in High-Energy Physics and Cosmology (CG 2003), Ft. Lauderdale, Florida, USA, 17-21 December, 2003.

[29] C.D. Froggatt, H.B. Nielsen and L.V. Laperashvili, Int. J. Mod. Phys. A**20**, 1268 (2005), hep-ph/0406110.

[30] H.B. Nielsen, Larisa Laperashvili, C.D. Froggatt, C.R. Das, *Top Yukawa New Strong Sector, Higgs Decay and Production*, a talk by H.B. Nielsen at the Corfu Conference, Greece, September, 1, 2016, PoS CORFU2015, 140 (2016), arXiv:1605.01169 [hep-ph].

[31] H.B. Nielsen, *F(750), We Miss You as Bound State of 6 Top and 6 Anti top*, arXiv:1610.00364.

[32] C.D. Froggatt and H.B. Nielsen, Phys. Lett. B**368**, 96 (1996), arXiv:hep-ph/9511371.

[33] M. Barriola and A. Vilenkin, *Gravitational Field of a Global Monopole*, Phys. Rev. Lett. **63**, 341 (1989).

[34] R. Basu, A. H. Guth and A. Vilenkin, *Quantum creation of topological defects during inflation*, Phys. Rev. D**44**, 340 (1991).

[35] A. Vilenkin, JCAP **1405**, 005 (2014), arXiv:1312.0682,

[36] J. Garriga, A. Vilenkin and Jun Zhang, JCAP **1602**, 064 (2016), arXiv:1512.01819.

[37] E.I. Guendelman and A. Rabinowitz, Phys. Rev. D**44**, 3152 (1991).

[38] F.B. Lustosa, M.E.X. Guimaraes, C.N. Ferreira and J.L. Neto, *Thermodynamical Analysis of a Black Hole with a Global Monopole Within a Class of a f(R) Gravity*, arXiv:1510.08176.

[39] O. Delice, JHEP **0311**, 058 (2003), arXiv:gr-qc/0307099.

[40] B.G. Sidharth, A. Das, C.R. Das, L.V. Laperashvili and H.B. Nielsen, Int. J. Mod. Phys. A**31**, 1630051 (2016), arXiv:1605.01169.

[41] C.R.Das, L.V. Laperashvili and A. Tureanu, Int. J. Mod. Phys. A**28**, 1350085 (2013), arXiv:1304.3069.

[42] G. t'Hooft, Nucl. Phys. B**79**, 276 (1974).

[43] A.M. Polyakov, Pisma Zh. Eksp. Teor. Fiz. **20**, 43 (1974) [JETP Lett. **20**, 194 (1974)].

[44] B.G. Sidharth, *The Chaotic Universe: From the Planck to the Hubble Scale* (Nova Science, New York, 2001).

[45] B.G. Sidharth, *The Universe of Fluctuations: The Architecture of Space-time and the Universe* (Springer, Netherlands, 2005).

[46] B.G. Sidharth, *The Thermodynamic Universe: Exploring the limits of physics* (World Scientific, Singapore, 2008).

[47] B.G. Sidharth, Found. Phys. **38** (1), 89 (2008); ibid., **38** (8), 695 (2008).

[48] B.G. Sidharth, Int. J. Th. Phys. **37**, 1307 (1998); ibid., **43** (9), 1857 (2004), arXiv:1201.0915.

[49] A. Das and B.G. Sidharth, EJTP **12**, No. IYL 15-34, 139 (2015).

[50] H.S. Snyder, Phys. Rev. **72**, 68 (1947).

[51] Ya.B. Zeldovich, JETP Lett. 6, 316 (1967).

[52] S. Weinberg, Phys. Rev. Lett. 43, 1566 (1979).

[53] A.A. Abrikosov, Soviet JETP **32**, 1442 (1957).

[54] H.B. Nielsen and P. Olesen, Nucl. Phys. B**61**, 45 (1973).

[55] B.G. Sidharth, A. Das and A.D. Roy, Int. J. Theor. Phys. **55**, 801 (2016).

[56] B.G. Sidharth, Nuovo. Cim. B**116**, 735 (2001), arXiv:physics/0106051.

[57] V.A. Belavin, M.N. Chernodub and I.E. Kozlov, Nucl. Phys. B**748**, 524 (2000), arXiv:hep-lat/0512030.

[58] M. N. Chernodub, Phys. Lett. **634**, 255 (2006), arXiv:hep-lat/0503018.

[59] C.D. Froggatt, L.V. Laperashvili and H.B. Nielsen, Phys. Atom. Nucl. **69**, 67 (2006) [Yad. Fiz. **69**, 3 (2006)], arXiv:hep-ph/0407102.

[60] G. Degrassi, S. Di Vita, J. Elias-Miro, J.R. Espinosa, G.F. Giudice, G. Isidori and A. Strumia, JHEP **1208**, 098 (2012), arXiv:1205.6497.

[61] D. Buttazzo, G. Degrassi, P.P. Giardino, G.F. Giudice, F. Salab, A. Salvio and A. Strumia, JHEP **1312**, 089 (2013), arXiv:1307.3536.



Fig. 1: Minima of the effective Higgs potential in the pure Standard Model, which correspond to the first Electroweak "true vacuum", and to the second Planck scale "false vacuum".



Fig. 2: The Wilson loop C (solid line) and its variation (dotted line).



Fig. 3: The RG evolution of the Higgs selfcoupling λ for $M_t \simeq 173.34$ GeV and $\alpha_s = 0.1184$ given by $\pm 3\sigma$. Blue lines present metastability for current experimental data, red (thick) line corresponds to the stability of the EW vacuum.



Fig. 4: The main Feynman diagram containing hedgehogs in the loop, which corrects the effective Higgs mass.